




Institute of Cosmophysical Research and Radio Wave Propagation FEB
RAS

Analysis of ionospheric parameters based on NARX neural networks


O.V. Mandrikova, Yu.A. Polozov

Time series structure of the critical frequency of the ionosphere


Ionospheric critical frequency time series for 2011 year (Paratunka station, Kamchatka, Russia)



foF2 time series for 2010 year (Paratunka station)



Space weather impact , 2017



<https://omniweb.gsfc.nasa.gov>


<https://cdaweb.gsfc.nasa.gov>

<http://isgi.unistra.fr>

<https://cdaweb.gsfc.nasa.gov>

<https://csu.ikir.ru/home>

Generalized recurrent network architecture



Input to hidden layer: $x(k), x(k-1), \dots, x(k-l_x), y(k), y(k-1), \dots, y(k-l_y)$

Delays relative to neural network output: $l_x = l_y$

Used time line delays:


$$l_x = l_y = 2$$

$$l_x = l_y = 5$$

$$l_x = l_y = 24$$


NARX SPA and NARX PA architectures

NARX SPA



$$y(k+1) = F[x(k), x(k-1), \dots, \\ x(k-l_x), \hat{y}(k), \\ \hat{y}(k-1), \dots, \hat{y}(k-l_y)].$$

NARX PA




$$y(k+1) = F[x(k), x(k-1), \dots, \\ x(k-l_x), y(k), \\ y(k-1), \dots, y(k-l_y)].$$


$F(\cdot)$ - neural network mapping function

Generating model data


Median foF2 (summer, low solar activity)




Noise



Train data



Model data NN errors




Standard deviations of neural network errors

Season	Time delay 2		Time delay 5		Time delay 24	
	NARX SPA	NARX PA	NARX SPA	NARX PA	NARX SPA	NARX PA
Winter (low solar activity)	0.47	0.57	0.42	0.53	0.31	0.3
Summer (low solar activity)	0.42	0.46	0.36	0.39	0.3	0.29
Winter (high solar activity)	0.5	0.78	0.43	0.67	0.3	0.29
Summer (high solar activity)	0.42	0.45	0.36	0.37	0.32	0.3

$$s = \sqrt{\frac{1}{N-1} \sum_{i=1}^N (e_i - \bar{e})^2},$$


$$\bar{e} = \frac{1}{N} \sum_{i=1}^N e_i$$

$$e_i = \hat{y}(i) - y(i)$$




Results of foF2 data processing for the period November 18 – 22, 2017

9



Results of foF2 data processing for the period August 03 – 08, 2018



Conclusions

NARX neural networks has shown that it can be used to detect anomalous features in the regular course of the foF2 time series. NN formed for different levels of solar activity and seasons. It demonstrated better approximation abilities for the NARX SPA architecture. With an increase of delay lines number, the ability to simulate a regular foF2 behavior increases. NARX neural networks can be used to detect anomalies in ionospheric parameters during periods of geomagnetic and ionospheric disturbances. The work was carried out as part of the implementation of the state task AAAA-A21-121011290003-0. The work was carried out by the means of the Common Use Center "North-Eastern Heliogeophysical Center" CKP_558279, USU 351757.

Mandrikova O., Fetisova N., Polozov Y. Hybrid Model for Time Series of Complex Structure with ARIMA Components // Mathematics. - 2021. - V.9. - P. 1122. <https://doi.org/10.3390/math9101122>.